

## ORIGINAL ARTICLE

*Sonographic Evaluation of Lens Thickness in Non – Diabetics and Diabetics with High-Resolution Ultrasound in a Tertiary Care Hospital*S. Aboorva<sup>1</sup>, K. A. Dharan Venkatesh<sup>1</sup>, V. Senthilnathan<sup>1</sup> and M. Krishna Kumar<sup>1</sup><sup>1</sup>Post Graduate Department of Radiology,<sup>1</sup>Trichy SRM Medical College and Research Centre, Trichy-621105, Tamil Nadu, India**Abstract:**

**Background:** Adult lenses have a thickness of approximately 4.0–5.0 mm and an average annual growth of 0.023 mm. It has been discovered that the human lens' thickness varies with age, sex, cataract type, and diabetes status. Myopia, presbyopia, and angle-closure glaucoma, among other disorders, are significantly influenced by the thickness of the crystalline lens. So, this study was undertaken to determine the lens thickness of diabetic individuals with a comparison of a control group. **Material and Methods:** An observational study with a control group was done at the Department of Radiology, to measure central lens thickness using High-resolution ultrasound of both eyes among Diabetic patients. They were compared to age, gender-matched healthy controls. The data were interpreted using MS Excel and SPSS software version 21. **Results:** The mean central lens thickness was  $3.759 \pm 0.294$  mm among Diabetic individuals and  $3.275 \pm 0.266$  mm among non-diabetic individuals in the right eye. The mean central lens thickness was  $3.746 \pm 0.432$  mm among Diabetic individuals and  $3.159 \pm 0.232$  mm among non-diabetic individuals in the left eye. Diabetic individuals had significantly thicker lenses on both sides compared with healthy individuals. Age is also a significant factor, which shows a positive correlation. **Conclusion:** The sonographic evaluation reveals the central lens thickness was increased among Diabetic eyes compared with non-diabetic eyes.

**Keywords:** Ultrasound, Lens thickness, Glaucoma, Diabetes, Cataract, Refraction

**Introduction:**

The human optical lens includes of a lens capsule,

epithelial cells, lens fibres and zonules. Adult lenses have a diameter of approximately 9.0–10.0 mm, a thickness of approximately 4.0–5.0 mm, and an average annual growth of 0.023 mm.<sup>[1]</sup> The lens geometry of Asian eyes can differ from Caucasian eyes due to the increased prevalence of myopia and extreme myopia in Asia.<sup>[2]</sup> Throughout life, the individual's crystalline lens tends to develop and produce new cells. The average human lens was 4 mm thick at age 20, 4.3 mm thick at age 40, 4.45 mm thick at age 50, 4.7 mm thick at age 60, and increased in thickness to beyond 4.7 mm after age 60.<sup>[3]</sup> It has been discovered that the human lens' thickness varies with age, sex, cataract type, and presence of Diabetes.<sup>[3]</sup> Myopia, presbyopia, and angle-closure glaucoma, among other disorders, are significantly influenced by the thickness of the crystalline lens.<sup>[4]</sup> The number of people having diabetes mellitus (DM) globally was anticipated by the International Diabetes Federation (IDF) to be 463 million in 2019 and to reach 700 million by 2045.<sup>[5]</sup> Diabetic retinopathy and papillopathy, cataract, glaucoma, and ocular surface disorders are the principal visual problems related to DM.<sup>[6]</sup> Diabetic retinopathy (DR), the most prevalent and distinct complication of DM, is a major risk factor for avoidable blindness in the adult working population.<sup>[5]</sup> DR affects one-third of diabetes individuals.<sup>[7]</sup> In 2020, there were thought to be 103.12 million adults worldwide who had DR; by 2045, that number is expected to increase to 160.50 million.<sup>[5]</sup> Diabetes can alter the lens in the following ways: Lens changes, refractive change, light scattering, chromophore accumulation, fluorescence, cataract and capsular changes.<sup>[8]</sup> The lens thickens with age, according to numerous investigations using ultrasound, Scheimpflug photography, partial coherence interferometry (PCI), and magnetic resonance imaging (MRI).<sup>[9]</sup> It has been established that USG's ocular biometry is more accurate and trustworthy than optical measurement.<sup>[10]</sup> Even in the presence of intervening optically opaque materials, ultrasound offers very cost-effective real-time cross-sectional images. In the 1950s,

Mundt and Hughes (A-scan) and Baum and Greenwood (B-scan) developed the first ultrasound systems for use in diagnostic imaging of the eye.<sup>[11]</sup> The standard technique for determining the biometric measurements of lens thickness is A-scan ultrasonography (A-scan US).<sup>[14]</sup> On the B-scan, the posterior portion displays normal, clear vitreous, and the anterior segment reveals the echo of the posterior lens capsule.<sup>[12]</sup> In comparison to traditional ophthalmic B-scanners, ultrasound biomicroscopy (UBM) employs frequencies that are substantially higher (35-50 MHz). As a result, ultrasound gets one step closer to fulfilling the requirements for the ideal imaging instrument, increasing resolution to 40  $\mu$ m or less. The cornea, iridocorneal angle, anterior chamber, iris, ciliary body, and lens are imaged with ultrasound biomicroscopy equipment, along with nearly all anterior segment anatomy and disease. In most cases, the crystalline lens produces specular surface reflectivity with little interior echoes.<sup>[11]</sup> Environmental variables and medical disorders can also affect refraction. As a result, diabetes mellitus may alter refraction both temporarily and permanently. It is widely believed that short-term fluctuations affect the lens' refraction mostly through changes in osmotic pressure brought on by variations in blood glucose levels.<sup>[13]</sup> The diabetes status also contributes to increasing lens thickness which affects vision. So, this study is an attempt to determine central lens thickness in diabetes compared with controls. The objectives of the study are to compare the central lens thickness of diabetic & non-diabetic patients (of similar age, and gender) using High Resolution Ultrasound and to compare the variation in central lens thickness between both eyes of diabetic patients.

### Material and Methods:

An observational study was done at Department of Radiology, Trichy SRM Medical College Hospital and Research Centre, Trichy for six months (July 2023 – January 2024) duration to determine the central lens thickness using high resolution ultrasound among patients with and without diabetes. Patients aged forty and years above referred for High-Resolution Ultrasound to the Department of Radiology, Trichy SRM Medical College Hospital and Research Centre, Trichy were selected as participants. Patients who had already undergone intraocular surgery/diagnosed with any corneal pathology, patients with inflammatory conditions like conjunctivitis, uveitis, scleritis, patients with associated hypertension and other co-morbidities, patients on statin medication and the participants not

willing to give consent were excluded from study. The cases (patients with diabetes) and controls (patient without diabetes) were selected using non probability sampling method. Based on inclusion and exclusion criteria eight-five cases and eighty-five controls were included in the study. Data was gathered in the Department of Radiology using semi-structured questionnaire among participants by interview method after getting their consent. Voluson S8, E6 machine with a multi-frequency linear transducer of 7.5 MHz was used for this study. Patients were placed in the supine posture for the B-Mode ultrasonography examination. Patients were scanned with their eyes closed while a minimum quantity of sonographic gel was applied to the probe. In the transverse projection, the probe was gently placed on the eyelid. Zooming in to fill the screen and reducing depth to include the back of the lens. The screen's centre was maintained for the lens. Midway between the front and posterior capsules of the lens, the maximum thickness of the lens was measured. Data was entered in Microsoft Excel 2019 and analysed using the software SPSS (Statistical Package of Social Sciences) version 21. The data were interpreted using frequencies (mean  $\pm$  SD) for quantitative variables and proportions (%) for qualitative variables. The independent T-test was utilised to test the association of central lens thickness among study and control groups. Approval was taken from the Institutional Ethical Committee of the hospital. Participants were informed about the study and their informed consent was obtained.

### Results:

This study was aimed to compare the central lens thickness of eight five diabetic and non-diabetic individuals to determine their central lens thickness. Table 1 describes the age group, gender and family history of diabetes of individuals. 15.3%, 32.9% and 51.8% of the diabetic individuals belonged to the age group of 41 – 50 years, 51 – 60 years and 61 – 80 years respectively. 31.8%, 38.8% and 29% of the non-diabetic individuals belonged to the age group of 41 – 50 years, 51 – 60 years, and 61 – 80 years respectively. Nearly half of the individuals who participated in this study from both groups were males. 42.4% of diabetic individuals had a family history of diabetes. Table 2 describes the association between demographic variables and diabetic status. Diabetes was more common in the age group 61 – 80 years. Diabetes was more common among individuals with a family history of diabetes. The mean age of

participants was  $64.047 \pm 9.71$  years in diabetic group and  $56.65 \pm 9.35$  years in non-diabetic group.

Table 1: Descriptive data of participants

Sr. No.	Variables		Diabetes (n = 85)	Non-Diabetes (n = 85)
1	Age group	41 – 50 years	13 (15.3%)	27 (31.8%)
		51 – 60 years	28 (32.9%)	33 (38.8%)
		61 – 80 years	44 (51.8%)	25 (29.4%)
2	Gender	Male	48 (56.5%)	49 (57.6%)
		Female	37 (43.5%)	36 (42.4%)
3	Family History of Diabetes	Present	36 (42.4%)	0
		Absent	49 (57.6%)	100 (100%)

Table 2: Association between demographic variables and diabetic status

Sr. No.	Variables		Diabetes (n = 85)	Non-Diabetes (n = 85)	p-value
1	Age group	41 – 50 years	13 (32.5%)	27 (67.5%)	0.001
		51 – 60 years	18 (35.3%)	33 (64.7%)	
		61 – 70 years	25 (59.5%)	17 (40.5%)	
		71 – 80 years	29 (78.4%)	8 (21.6%)	
2	Gender	Male	48 (49.5%)	49 (50.5%)	0.877
		Female	37	36	

			(50.7%)	(49.3%)	
3	Family History of Diabetes	Present	36 (100%)	0	0.001
		Absent	49 (36.6%)	85 (63.4%)	

Table 3 shows central lens thickness among individuals. The mean central lens thickness was found to be higher in diabetic individuals compared with non-diabetic individuals and this seems to be statistically significant. The mean central lens thickness was  $3.759 \pm 0.294$  mm (ranging from 3 – 4.3mm) among diabetic individuals and  $3.275 \pm 0.266$  mm (ranging from 3 – 4 mm) among non-diabetic individuals in the right eye. The mean central lens thickness was  $3.746 \pm 0.432$  mm among diabetic individuals and  $3.159 \pm 0.232$  mm among non-diabetic individuals in the left eye.

Table 3: Central Lens thickness among individuals (in mm)

Sr. No.	Variable	Diabetes	Non-Diabetes	p-value
1	Right	$3.759 \pm 0.294$	$3.275 \pm 0.266$	0.001
2	Left	$3.746 \pm 0.432$	$3.159 \pm 0.232$	0.001

Table 4: Correlation of age with central lens thickness

Sr. No.	Variable	Diabetes	Non-Diabetes
1	Right	$r = 0.537$ $p = <0.001$	$r = 0.585$ $p = <0.001$
2	Left	$r = 0.442$ $p = <0.001$	$r = 0.522$ $p = <0.001$

The mean central lens thickness was  $3.758 \pm 0.294$  mm and  $3.746 \pm 0.432$  mm in the right eye and left eye of diabetes patients respectively where there was no significant difference in the right and left eye ( $p = 0.820$ ). Table 4 shows the correlation of age with central lens thickness. Age shows a positive correlation ( $r = 0.537$ ) in the right eye and the left eye ( $r = 0.517$ ) with a statistically significant p-value (p-value = 0.001). Therefore, it is seen that the central lens thickness increases with respect to age. Age showed a positive correlation with central lens thickness and was found to

be statistically significant. Table 5 shows the association of central lens thickness with gender variables. There was no significant change in the central lens thickness of both eyes among male and female individuals.

Table 5: Association of central lens thickness with gender variable (in mm)

Sr. No.	Group	Variable	Male	Female	p-value
1	Diabetes	Right	3.783±0.250	3.727±0.343	0.385
		Left	3.792±0.501	3.686±0.320	0.269
2	Non-Diabetes	Right	3.294±0.305	3.25±0.204	0.457
		Left	3.171±0.27	3.142±0.169	0.562

### Discussion:

The mean central lens thickness of the right eye was  $3.759 \pm 0.294$  mm among Diabetic and  $3.275 \pm 0.266$  mm among non-diabetic. The mean central lens thickness of the left eye was  $3.746 \pm 0.432$  mm among diabetic and  $3.159 \pm 0.232$  mm among non-diabetic. Deepa KB et al<sup>[14]</sup> in a study found that lens thickness of the right eye was  $4.1 \pm 2.9$  mm among diabetic retinopathy patients, while it was  $3.7 \pm 1.3$  mm among normal healthy individuals, and the difference among groups was found to be statistically significant. The difference in lens thickness of the left eye was found to be statistically significant among DR cases ( $4.2 \pm 0.3$  mm) and normal controls ( $3.7 \pm 0.1$  mm). According to Shrestha S et al<sup>[15]</sup> in Nepal, diabetic participants' crystalline lenses ( $4.33 \pm 0.38$  mm) were substantially thicker than those of the control group ( $4.05 \pm 0.55$  mm). Studies found that diabetes contributes to lens thickness significantly, which was similar to our study results. Thomas et al<sup>[16]</sup> among the Nigerian population found that Diabetes ( $3.76 \pm 0.45$  mm) had thicker lenses compared non diabetes ( $3.56 \pm 0.34$  mm). Ye L et al<sup>[17]</sup> among Chinese adults aged fifty and above found that Lens thickness among Diabetics ( $4.63 \pm 0.32$  mm) was significantly different compared to the control group

( $4.69 \pm 0.36$  mm). Thomas et al<sup>[16]</sup>, Meng et al<sup>[2]</sup>, Sharma et al<sup>[18]</sup>, and Praveen MR et al<sup>[3]</sup> found that increasing lens thickness due to increasing age seems to be a positive correlation between age and lens thickness. Praveen MR et al<sup>[3]</sup> concluded that with each decade of age advancement, lens thickness was increased by 0.155 mm. Pierro L et al<sup>[19]</sup> found that there significant changes in lens thickness among Diabetes compared with Non Diabetes among patients with no and proliferative retinopathy and also the author stated that lens thickness had significant correlation with duration of diabetes. Diabetics have thicker lenses than healthy patients, primarily because of cortical thickening. The lens's front and rear curvatures getting steeper, and the anterior chamber getting shallower, are other biometric alterations. After correcting for age, insulin-dependent (type 1) diabetics exhibit the most substantial alterations, and the duration of diabetes has a significant independent impact on lens thickness. This rise among type 1 diabetics does appear to be influenced by the prevalence of diabetic retinopathy.<sup>[8]</sup> Although this study does not stratify type of diabetes, lens thickness has significantly increased among diabetic individuals.

### Conclusion:

The diabetic lens is larger than average, more susceptible to refractive alteration, and more likely to develop a cataract, sometimes of a particular variety. As a result, reliable lens thickness measurements are crucial for research on the onset of refractive error, cataract surgery, and angle closure glaucoma. Therefore, regular ocular follow-up with high-resolution ultrasound is necessary to aid in the early diagnosis of lenticular thickness changes that may increase the risk of cataract formation, and reduce vision, and glaucomatous changes that may have preventive implications.

### Limitations:

The sample size was small for generalising the results. The variables like associated co-morbid conditions, duration of diabetes, and type of diabetes were not included.

Sources of supports: Nil

Conflicts of Interest: Nil

### References

1. Ruan X, Liu Z, Luo L, Liu Y. Structure of the lens and its associations with the visual quality. *BMJ Open Ophthalmology* 2020 Sep 1;5(1):e000459.
2. Meng J, Wei L, He W, Qi J, Lu Y, Zhu X. Lens thickness and associated ocular biometric factors among cataract patients in Shanghai. *Eye and Vision* 2021 May 31;8(1):22.



3. Praveen MR, Vasavada AR, Shah SK, Shah CB, Patel UP, Dixit NV, et al. Lens thickness of Indian eyes: impact of isolated lens opacity, age, axial length, and influence on anterior chamber depth. *Eye* 2009 Jul;23(7):1542–1548.
4. Hamzeh N, Moghimi S, Latifi G, Mohammadi M, Khatibi N, Lin SC. Lens thickness assessment: anterior segment optical coherence tomography versus A-scan ultrasonography. *International Journal of Ophthalmology* 2015 Dec 18;8(6):1151–1155.
5. Teo ZL, Tham YC, Yu M, Chee ML, Rim TH, Cheung N, et al. Global Prevalence of Diabetic Retinopathy and Projection of Burden through 2045: Systematic Review and Meta-analysis. *American academy of Ophthalmology* 2021 Nov;128(11):1580–1591.
6. Sayin N, Kara N, Pekel G. Ocular complications of diabetes mellitus. *World Journal of Diabetes* 2015 Feb 15;6(1):92–108.
7. Hashemi H, Rezvani F, Pakzad R, Ansari-pour A, Heydarian S, Yekta A, et al. Global and Regional Prevalence of Diabetic Retinopathy; A Comprehensive Systematic Review and Meta-analysis. *Seminars Ophthalmology* 2022 Apr 3;37(3):291–306.
8. Bron AJ, Sparrow J, Brown NA, Harding JJ, Blakely R. The lens in diabetes. *Eye (Lond)* 1993;7 (2):260–275.
9. Richdale K, Bullimore MA, Zadnik K. Lens Thickness with Age and Accommodation by Optical Coherence Tomography. *Ophthalmic and Physiological Optics* 2008; Sep;28(5):441–447.
10. Zeng Y, Liu Y, Liu X, Chen C, Xia Y, Lu M, et al. Comparison of lens thickness measurements using the anterior segment optical coherence tomography and A-scan ultrasonography. *Investigate Ophthalmology and Visual Science* 2009 Jan;50(1):290–294.
11. Silverman RH. High-resolution ultrasound imaging of the eye – a review. *Clinical and Experimental Ophthalmology* 2009 Jan; 37 (1) : 54–67.
12. Aironi V, Gandage S. Pictorial essay: B-scan ultrasonography in ocular abnormalities. *The Indian Journal of Radiology & Imaging* 2009 Apr 30;19:109–115.
13. Løgstrup N, Sjølie AK, Kyvik KO, Green A. Lens thickness and insulin dependent diabetes mellitus: a population based twin study. *British Journal of Ophthalmology* 1996 May 1;80(5):405–408.
14. Deepa KP, Ghafoor N, Haque Z, Islam U, Jubaida R. Comparison of Ocular Lens Thickness of Type-2 Diabetic Retinopathy Patients with Non-diabetic Control, Measured by High Resolution Ultrasonography. *Ibrahim Cardiac Medical Journal* 10((1 & 2)):40–44.
15. Shrestha S, Kaini KR. The Influence of Diabetes Mellitus on Lenticular Thickness. *American Journal of Public Health Research* 2015 Oct 28;3(5A):91–94.
16. Thomas AA, Ayoola OO, Ikem R, Onakpoya TH, Dada AS, Bello T. Sonographic Evaluation of Lens Thickness in Diabetes Mellitus in a Nigerian Tertiary Hospital. *Nigerian Journal of Ophthalmology* 2018 Dec;26(2):104.
17. Ye L, He J, Zhang X, Xu Y, Chen Q, Yin Y, et al. The associations of lens power with age, axial length and type 2 diabetes mellitus in Chinese adults aged 50 and above. *Eye Vision (Lond)* 2020 Dec 1;7(1):57.
18. Sharma S, Sharma AK, Gupta S, Gupta D. Lens thickness and associated ocular biometric factors among cataract patients in tertiary hospital. *International Journal of Research in Medical Sciences* 2023 Feb 7;11(3):852–856.
19. Pierro L, Brancato R, Zaganelli E, Guarisco L, Calori G. Correlation of lens thickness with blood glucose control in diabetes mellitus. *Acta Ophthalmologica Scandinavica* 1996 Dec;74(6):539–541.

**Address for correspondence:**

Dr. K. A. Dharan Venkatesh  
Associate professor of Radiology  
Trichy SRM Medical College Hospital and Research  
Center, Trichy- 621105, Tamil Nadu, India  
Mobile no: +91 9944369232  
Email: dr.dharanvenkat@gmail.com

**How to cite this article:**

S. Aboorva, K. A. Dharan Venkatesh, V. Senthilnathan and R. Vinnarasi. Sonographic Evaluation of Lens Thickness in Non – Diabetics and Diabetics with High-Resolution Ultrasound in a Tertiary Care Hospital. Walawalkar International Medical Journal 2024;11(1):30-36  
<http://www.wimjournal.com>.

Received date:11/03/2024

Revised date:22/07/2024

Accepted date: 23/07/2024